



proximal 18q deletion syndrome

Proximal 18q deletion syndrome is a chromosomal condition that occurs when a piece of the long (q) arm of chromosome 18 is missing. The term "proximal" means that the missing piece occurs near the center of the chromosome. Individuals with proximal 18q deletion syndrome have a wide variety of signs and symptoms. Because only a small number of people are known to have this type of deletion, it can be difficult to determine which features should be considered characteristic of the disorder.

Most people with proximal 18q deletion syndrome have delayed development of skills such as sitting, crawling, walking, and speaking, and intellectual disability that can range from mild to severe. In particular, vocabulary and the production of speech (expressive language skills) may be delayed. Recurrent seizures (epilepsy) and weak muscle tone (hypotonia) often occur in this disorder. Affected individuals also frequently have behavioral problems such as hyperactivity, aggression, excessive eating (hyperphagia) that can lead to obesity, and features of autism spectrum disorders that affect communication and social interaction. Some affected individuals have an unusually large head size (macrocephaly).

People with proximal 18q deletion syndrome often have characteristic facial features including a prominent forehead, droopy eyelids (ptosis), short and slightly downslanting openings of the eyes (palpebral fissures), puffy tissue around the eyes (periorbital tissue), and full cheeks.

Frequency

Deletions from the q arm of chromosome 18 occur in an estimated 1 in 40,000 newborns worldwide. However, only a small number of these individuals have deletions in the region associated with proximal 18q deletion syndrome. At least 15 people with proximal 18q deletion syndrome have been described in the medical literature.

Genetic Changes

Proximal 18q deletion syndrome is caused by a deletion of genetic material from one copy of chromosome 18. The deletion occurs near the middle of the long arm of the chromosome, typically in an area between regions called 18q11.2 and 18q21.2. The size of the deletion varies among affected individuals. The signs and symptoms of proximal 18q deletion syndrome are thought to be related to the loss of multiple genes from this part of chromosome 18. Researchers are working to determine how the loss of specific genes in this region contributes to the various features of this disorder.

Inheritance Pattern

Proximal 18q deletion syndrome is considered to be an autosomal dominant condition. This means that a deletion in one of the two copies of chromosome 18 in each cell is sufficient to cause the disorder's characteristic features.

Most cases of proximal 18q deletion syndrome are the result of a new (de novo) deletion and are not inherited from a parent. The deletion occurs most often as a random event during the formation of reproductive cells (eggs or sperm) or in early fetal development. Affected people typically have no history of the disorder in their family.

Other Names for This Condition

- 18q deletion syndrome
- 18q- syndrome
- chromosome 18 deletion syndrome
- chromosome 18 long arm deletion syndrome
- chromosome 18q monosomy
- chromosome 18q- syndrome
- del(18q) syndrome
- monosomy 18q

Diagnosis & Management

Genetic Testing

- Genetic Testing Registry: Deletion of long arm of chromosome 18
<https://www.ncbi.nlm.nih.gov/gtr/conditions/C0432443/>

Other Diagnosis and Management Resources

- University of Texas Health Science Center: Chromosome 18 Clinical Research Center
<http://www.pediatrics.uthscsa.edu/centers/chromosome18/>

General Information from MedlinePlus

- Diagnostic Tests
<https://medlineplus.gov/diagnostictests.html>
- Drug Therapy
<https://medlineplus.gov/drugtherapy.html>
- Genetic Counseling
<https://medlineplus.gov/geneticcounseling.html>

- Palliative Care
<https://medlineplus.gov/palliativecare.html>
- Surgery and Rehabilitation
<https://medlineplus.gov/surgeryandrehabilitation.html>

Additional Information & Resources

MedlinePlus

- Health Topic: Developmental Disabilities
<https://medlineplus.gov/developmentaldisabilities.html>
- Health Topic: Epilepsy
<https://medlineplus.gov/epilepsy.html>

Genetic and Rare Diseases Information Center

- Proximal chromosome 18q deletion syndrome
<https://rarediseases.info.nih.gov/diseases/10866/proximal-chromosome-18q-deletion-syndrome>

Educational Resources

- Merck Manual Consumer Version: Overview of Chromosomal Disorders
<http://www.merckmanuals.com/home/children-s-health-issues/chromosomal-and-genetic-abnormalities/overview-of-chromosomal-disorders>
- Unique: 18q Deletions from 18q11.2 to 18q21.2
<http://www.rarechromo.org/information/Chromosome%2018/18q%20deletions%20from%20%2018q11.2%20to%2018q21.2%20FTNW.pdf>

Patient Support and Advocacy Resources

- Chromosome Disorder Outreach
<http://chromodisorder.org/>
- Epilepsy Foundation
<http://www.epilepsy.com/>
- March of Dimes: Chromosomal Conditions
<http://www.marchofdimes.org/baby/chromosomal-conditions.aspx>
- National Organization for Rare Disorders
<https://rarediseases.org/rare-diseases/chromosome-18q-syndrome/>
- The Arc: for People with Intellectual and Developmental Disabilities
<http://www.thearc.org/>

- The Chromosome 18 Registry and Research Society
<https://www.chromosome18.org/18q/proximal-18q/>
- Unique: The Rare Chromosome Disorder Support Group
<http://www.rarechromo.org/html/home.asp>

Scientific Articles on PubMed

- PubMed
<https://www.ncbi.nlm.nih.gov/pubmed?term=%28proximal+18q+deletion+syndrome%29+AND+english%5Bla%5D+AND+human%5Bmh%5D>

OMIM

- CHROMOSOME 18q DELETION SYNDROME
<http://omim.org/entry/601808>

Sources for This Summary

- Buysse K, Menten B, Oostra A, Tavernier S, Mortier GR, Speleman F. Delineation of a critical region on chromosome 18 for the del(18)(q12.2q21.1) syndrome. *Am J Med Genet A*. 2008 May 15; 146A(10):1330-4. doi: 10.1002/ajmg.a.32267.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/18412119>
- Cody JD, Ghidoni PD, DuPont BR, Hale DE, Hilsenbeck SG, Stratton RF, Hoffman DS, Muller S, Schaub RL, Leach RJ, Kaye CI. Congenital anomalies and anthropometry of 42 individuals with deletions of chromosome 18q. *Am J Med Genet*. 1999 Aug 27;85(5):455-62.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/10405442>
- Cody JD, Sebold C, Heard P, Carter E, Soileau B, Hasi-Zogaj M, Hill A, Rupert D, Perry B, O'Donnell L, Gelfond J, Lancaster J, Fox PT, Hale DE. Consequences of chromosome 18q deletions. *Am J Med Genet C Semin Med Genet*. 2015 Sep;169(3):265-80. doi: 10.1002/ajmg.c.31446.
Review.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/26235940>
- Cody JD, Sebold C, Malik A, Heard P, Carter E, Crandall A, Soileau B, Semrud-Clikeman M, Cody CM, Hardies LJ, Li J, Lancaster J, Fox PT, Stratton RF, Perry B, Hale DE. Recurrent interstitial deletions of proximal 18q: a new syndrome involving expressive speech delay. *Am J Med Genet A*. 2007 Jun 1;143A(11):1181-90.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/17486614>
- Feenstra I, Vissers LE, Orsel M, van Kessel AG, Brunner HG, Veltman JA, van Ravenswaaij-Arts CM. Genotype-phenotype mapping of chromosome 18q deletions by high-resolution array CGH: an update of the phenotypic map. *Am J Med Genet A*. 2007 Aug 15;143A(16):1858-67.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/17632778>
- Hale DE, Cody JD, Baillargeon J, Schaub R, Danney MM, Leach RJ. The spectrum of growth abnormalities in children with 18q deletions. *J Clin Endocrinol Metab*. 2000 Dec;85(12):4450-4.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/11134092>
- Imataka G, Ohwada Y, Shimura N, Yoshihara S, Arisaka O. Del(18)(q12.2q21.1) syndrome: a case report and clinical review of the literature. *Eur Rev Med Pharmacol Sci*. 2015 Sep;19(17):3241-5.
Review.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/26400529>

- Lancaster JL, Cody JD, Andrews T, Hardies LJ, Hale DE, Fox PT. Myelination in children with partial deletions of chromosome 18q. *AJNR Am J Neuroradiol*. 2005 Mar;26(3):447-54.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/15760848>
 - Schaub RL, Hale DE, Rose SR, Leach RJ, Cody JD. The spectrum of thyroid abnormalities in individuals with 18q deletions. *J Clin Endocrinol Metab*. 2005 Apr;90(4):2259-63.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/15671099>
 - Tinkle BT, Christianson CA, Schorry EK, Webb T, Hopkin RJ. Long-term survival in a patient with del(18)(q12.2q21.1). *Am J Med Genet A*. 2003 May 15;119A(1):66-70.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/12707962>
-

Reprinted from Genetics Home Reference:

<https://ghr.nlm.nih.gov/condition/proximal-18q-deletion-syndrome>

Reviewed: February 2017

Published: March 21, 2017

Lister Hill National Center for Biomedical Communications
U.S. National Library of Medicine
National Institutes of Health
Department of Health & Human Services